

CAPOX + Bevacizumab

Your chemotherapy treatment is called CAPOX + bevacizumab. It is commonly used to treat colon and rectal cancer and has also been used to treat other diseases. CAPOX + bevacizumab is made up of three drugs:

- capecitabine (kape-SITE-a-been) or Xeloda® (zeh-LOE-duh)
- oxaliplatin (ox-AL-i-pla-tin) or Eloxatin® (ee-LOX-a-tin)
- bevacizumab (be-va-SIZ-you-mab) or Avastin® (ah-VA-stin)

Capecitabine and oxaliplatin prevent cancer cells from dividing and growing, and can eventually cause the cancer cells to shrink and die. Bevacizumab is a new type of drug, called a monoclonal antibody, which targets cancer cells more precisely than chemotherapy drugs.

What Do I Need to Know Before Starting Treatment?

Be sure to tell your healthcare provider about any prescription or over-the-counter products you are taking, including dietary supplements, vitamins, herbal medicines and homeopathic remedies.

Use an effective birth control method while you are taking these drugs. Chemotherapy drugs can cause harm to a fetus, so be sure to tell your healthcare provider right away if you or your partner become pregnant.

Avoid breastfeeding during treatment. It is not known if these drugs pass into breast milk.

Some chemotherapy drugs can cause sterility. Talk with your healthcare provider about your options if you want to have children in the future.

Do not get any immunizations or vaccinations while taking chemotherapy drugs without the approval of your healthcare provider.

What Do I Need to Know Before Starting CAPOX + Bevacizumab?

Capecitabine can increase the effects of blood thinners such as warfarin (Coumadin®). It is extremely important to tell your healthcare provider if you are taking warfarin or another blood thinner so that your healthcare provider can check your blood's ability to clot. The dose of your blood thinner may have to be decreased while you are taking capecitabine.

Capecitabine can cause severe diarrhea. See "What are the Possible Side Effects" for how to manage diarrhea. If your diarrhea is severe, call your healthcare provider.

Capecitabine and bevacizumab can cause heart problems such as irregular heartbeat, angina, electrocardiographic (EKG) changes, heart attack and heart failure. These side effects may be more common in patients with a history of heart disease. If you have any heart problems, call your healthcare provider.

Oxaliplatin commonly causes some nerve cells to stop functioning properly. This condition is called neuropathy (see "What Are the Possible Side Effects?"). Oxaliplatin can cause two types of neuropathy:

- Early onset neuropathy starts within a few hours to two days after starting treatment and usually goes away within 14 days. Exposure to cold temperatures and cold objects can cause symptoms or make symptoms worse. Avoid cold weather, cold foods and cold drinks for five days after treatment.
- Oxaliplatin can cause chronic neuropathy that can last for weeks or months. Chronic neuropathy usually starts after several treatments with oxaliplatin.

A less common side effect of oxaliplatin-related neuropathy is called pharyngolaryngeal dysesthesia. This is the sensation of tightness or discomfort in the throat, which can make it feel difficult to breathe or swallow. Although this symptom may be frightening, it is just a sensation and does not interfere with breathing. The sensation usually goes away on its own after a few minutes.

Oxaliplatin can cause irritation if it accidentally leaks out of the vein. Tell your healthcare provider right away if you have redness, pain, burning or swelling around the injection site during your treatment.

Oxaliplatin can damage the kidneys. Tell your healthcare provider if you have a history of kidney disease.

Oxaliplatin can cause a severe allergic reaction a few minutes after treatment. Tell your healthcare provider right away if you develop a rash, swollen lips or tongue or if you have trouble breathing.

In rare cases, oxaliplatin can cause a lung condition called pulmonary fibrosis. Tell your healthcare provider immediately if you suddenly develop a cough or are short of breath,

In rare cases, the allergic reactions and pulmonary fibrosis caused by oxaliplatin can be life-threatening.

In rare cases, bevacizumab can cause severe infusion reactions, such as trouble breathing, very high blood pressure, wheezing, chest pain, headache, chills and sweating. If this reaction occurs, it is usually with the first dose.

Bevacizumab can cause gastrointestinal perforation (holes in the stomach, intestines or colon) that must be repaired through surgery. Call your healthcare provider right away if you have abdominal pain with nausea, vomiting, blood in your stool, constipation or fever.

Bevacizumab can cause slow or incomplete wound healing. Tell your healthcare provider if you have wounds that do not heal. You should not have any elective surgery while being treated with bevacizumab.

Mild bleeding, such as nosebleeds, is common with bevacizumab. If you have a nosebleed, sit with your head tipped slightly forward and pinch the bridge of your nose. Call your healthcare provider if you feel dizzy or faint, or if the bleeding doesn't stop after 10 to 15 minutes.

Bevacizumab may cause a severe increase in blood pressure. Your healthcare provider will check your blood pressure regularly and may give you medicine to treat high blood pressure.

Bevacizumab can cause too much protein in the urine, which can lead to kidney damage. Your healthcare provider will check your urine regularly for signs of kidney damage.

In rare cases, bevacizumab can cause serious blood clots that can result in stroke. Older adults may be most likely to develop blood clots. Tell your healthcare provider if you have a history of heart disease.

Your treatment can interfere with other medicines, including:

- Warfarin (Coumadin®)
- Phenytoin (Dilantin®) – if you are taking this medication, your blood levels will need to be checked regularly
- Folic acid
- Drugs that cause harm to the kidney, such as amphotericin B and aminoglycosides (such as tobramycin, gentamicin or amikacin)

Please note this list is a summary and does not contain all possible drug interactions. Contact your healthcare provider if you are taking any medications that can interact with your treatment.

You should not take this treatment if you:

- Are allergic to capecitabine, oxaliplatin, other platinum agents, bevacizumab or any components of these drugs or if you are allergic to 5-fluorouracil (5FU) because capecitabine is converted to 5FU in the body.
- Have a condition known as dihydropyrimidine dehydrogenase (DPD) deficiency.

How Is the Treatment Given?

Your healthcare provider will give you oxaliplatin and bevacizumab by injection into a vein. The doses you receive will be based on your weight and height. Your healthcare provider will determine the number of treatments you receive.

Capecitabine is usually taken twice a day, in the morning and evening, for 14 days, followed by a seven-day rest period. Follow your healthcare provider's directions if you are given a different dosing schedule.

Capecitabine should be swallowed with water and taken within 30 minutes after a meal. If you miss a dose, do not take the missed dose or double the next dose. Instead, continue your regular dosing schedule and check with your healthcare provider. If you are given two different strengths of capecitabine, make sure that you can tell the difference between the 150 mg. (light peach with "150" on one side) and the 500 mg. (darker peach with "500" on one side) tablets.

Store capecitabine at room temperature away from children and pets. If you take too much capecitabine, contact your healthcare provider, local poison control center or emergency room right away.

You may be given medicines to help prevent and control nausea and vomiting before you receive your treatment. These medicines may be given either by mouth or by injection into a vein.

If you are given any medicine to take at home, do not share it with others. Sharing this medication with anyone else could be harmful.

When Should I Call My Healthcare Provider?

Call your healthcare provider right away if you have any of the following symptoms:

- Shaking chills or fever of 100.5 degrees F or higher
- Unusual bleeding, easy bruising or pinpoint red spots on your skin
- Vomiting that is severe or lasts several hours
- Painful or frequent urination or blood in your urine
- Diarrhea that causes an additional four bowel movements a day, diarrhea that lasts more than one day, diarrhea at night or diarrhea with fever, cramps or bloody stools
- Irregular or rapid heart beat, chest pain, chest tightness or shortness of breath
- Dizziness or feeling lightheaded
- Inability to eat or weight loss

What Are the Possible Side Effects?

All drugs can cause side effects, but every person reacts differently to each drug. The following chart lists the possible side effects that can occur with your treatment, how to recognize and minimize symptoms and possible treatments. The side effects are grouped by how often the side effect occurs: Common (occurs in more than 25 percent of patients), Less Common (occurs in 5 to 25 percent of patients) or Rare (occurs in less than 5 percent of patients).

Side Effect	How to Minimize Side Effect	Possible Treatments
<p>Nausea/Vomiting (Common. Symptoms are generally mild to moderate.)</p> <ul style="list-style-type: none"> • Feeling queasy or sick to your stomach 	<ul style="list-style-type: none"> • Eat small, frequent meals and bland foods—such as bananas, rice, applesauce and toast. • Eat food cold or at room temperature so the smell of food will not bother you. • Avoid fried, spicy or fatty foods. • Eat and drink slowly. • Drink plenty of liquids during the day, but to avoid bloating, drink small amounts of liquid during meals. 	<ul style="list-style-type: none"> • You will be given medicine to help reduce nausea and vomiting.
<p>Diarrhea (Common. Symptoms are generally mild to moderate.)</p> <ul style="list-style-type: none"> • Loose or watery stools several times a day • Abdominal cramping, gas and bloating 	<ul style="list-style-type: none"> • Eat small, frequent meals and bland foods—such as bananas, rice, applesauce and toast. • Avoid caffeine; alcohol; raw fruits and vegetables; raw eggs; undercooked meats; spicy, fatty and greasy foods; milk and dairy products; foods that cause gas, such as beans and other legumes; high fiber and high-fat foods; foods left un-refrigerated for more than two hours (one hour for egg dishes and cream or mayonnaise-based foods); bulk laxatives; and stool softeners. 	<ul style="list-style-type: none"> • Drink eight to ten glasses of clear liquids every day. • Your healthcare provider may prescribe medicine to help treat diarrhea.
<p>Hand-Foot Syndrome (Common. Symptoms are generally mild to moderate.)</p> <ul style="list-style-type: none"> • Swelling or redness in the hands and feet that can prevent normal activity • Cracked or peeling skin 	<ul style="list-style-type: none"> • Wear gloves to wash dishes. • Avoid using harsh household cleaners. • Avoid prolonged exposure to heat. • Use creams or moisturizers regularly. Try wearing cotton gloves on your hands. • Avoid using perfume, cologne or aftershave since these products can be irritating to the 	<ul style="list-style-type: none"> • Your healthcare provider may prescribe a cream to help with skin reactions. • Your healthcare provider may need to decrease the dose or delay further chemotherapy.

Side Effect	How to Minimize Side Effect	Possible Treatments
<ul style="list-style-type: none"> • Tingling, numbness or pain in the hands or soles of the feet 	<p>skin.</p>	
<p>Neuropathy (Common. Symptoms are generally mild to moderate.)</p> <ul style="list-style-type: none"> • Numbness or tingling feeling in the hands or feet • Muscle cramps • Loss of balance • Difficulty buttoning buttons or picking up objects • Decreased awareness of heat or cold in fingertips and toes • Difficulty hearing 	<ul style="list-style-type: none"> • Try to avoid extreme heat. • Wear mittens or gloves, socks and scarves. • If your fingers are numb, be careful with sharp objects. • Beware of hot coffee mugs, pots and pans and dishwasher—you may not feel the heat until you are burned. • If you feel unsteady, be careful on stairs and in the shower. • Avoid exposure to cold, including cold foods and drinks, for five days after treatment. 	<ul style="list-style-type: none"> • Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.
<p>Alopecia or Hair Loss (Less Common. Does not usually cause complete hair loss.)</p> <ul style="list-style-type: none"> • More than normal amount of hair loss in your brush, in the shower or on your pillow after sleeping • Loss of body hair 	<p>Alopecia cannot be prevented but here are tips to help with hair loss:</p> <ul style="list-style-type: none"> • Use a soft hairbrush. Do not use brush rollers, color treat your hair or get a permanent. • Avoid daily hair washing, use a mild shampoo and avoid using a hairdryer, or use a low setting if you must use one. • Have your hair cut short; this will make it look fuller. • Your insurance might cover a wig. Ask your healthcare provider for a prescription for a "hair prosthesis". Your hair color and style can be better matched if you shop for a wig before losing a lot of hair. • Use sunscreen or wear a hat or scarf to protect your scalp from the sun. 	
<p>Anorexia or Appetite Loss (Less Common)</p> <ul style="list-style-type: none"> • Not having an appetite • Feeling too nauseated to eat • Metallic or medicinal taste • Change in taste causing dislike for certain foods 	<ul style="list-style-type: none"> • Try eating six to eight small meals or snacks each day instead of three larger meals. • Vary your diet and try new foods and recipes. • Take a walk before meals, when possible. This may make you feel hungrier. • Eat with friends or family. When eating alone, listen to the radio or watch TV. 	<ul style="list-style-type: none"> • Cook dinners ahead of time and freeze them in small portions so that cooking smells are minimized. • Let others help with food, but ask that foods be prepared in small portions that can be frozen. And don't hesitate to let them know which foods to avoid. • Add mild spices to change flavor. • It might be helpful to have a program, such as Meals on Wheels, deliver food to you.
<p>Mouth Sores and Pain (Less Common. Symptoms are generally mild.)</p> <ul style="list-style-type: none"> • Pain, swelling or redness of the mouth, tongue and throat • "Coated tongue" • Difficulty talking, swallowing or eating • Bleeding ulcers and infection 	<ul style="list-style-type: none"> • Brush teeth two to four times a day using a soft bristle brush and fluoride toothpaste. • Use non-waxed dental floss daily. • Ask your healthcare provider to recommend a mouthwash that does not contain alcohol. • Sip water during the day and use sugar-free candy or gum to keep your mouth wet. • Eat food cold or at room temperature. • Eat soft or pureed food. • Avoid food that is acidic, spicy, salty, dry or rough, such as toast. 	<ul style="list-style-type: none"> • You may be given medicine to help treat pain. • You may be given medicine to treat fungal or viral infections.
<p>Risk of Infection (Less Common)</p> <ul style="list-style-type: none"> • Fever and chills • Painful urination • Sore throat and cough • Nasal congestion • Swelling or redness of the skin at the site of a wound 	<ul style="list-style-type: none"> • Wash your hands often. • Brush and floss your teeth daily. • Clean cuts right away with warm water, soap and antiseptic. • When your white blood cell count is low, stay away from crowds and people with colds or other illnesses. 	<ul style="list-style-type: none"> • You may be given medicine to increase your white blood cell count. • You may be given an antibiotic to treat or prevent infection. • Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.
<p>Bleeding (Less Common)</p> <ul style="list-style-type: none"> • Unusual bleeding or easy bruising • Black or tar-like stools • Blood in your urine • Pinpoint red spots on your skin 	<ul style="list-style-type: none"> • Avoid aspirin and aspirin-like drugs, such as ibuprofen. • Use caution with sharp objects, such as razors and nail cutters. • Avoid activities that can cause cuts, bumps 	<ul style="list-style-type: none"> • You may be given medicine to increase your platelet count. • Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.

Side Effect	How to Minimize Side Effect	Possible Treatments
<ul style="list-style-type: none"> Bleeding gums or nosebleeds 	and bruises.	
Anemia (Rare) <ul style="list-style-type: none"> Fatigue or weakness Dizziness Pale skin Feeling out of breath Feeling cold 	<ul style="list-style-type: none"> Plan rest periods throughout the day. Organize daily activities so that you conserve your energy. Try to eat a well balanced diet and drink plenty of fluids. Stand up slowly to avoid getting dizzy. 	<ul style="list-style-type: none"> You may be given medicine to increase your red blood cell count. Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.

What Are The Other Possible Side Effects?

The chart below lists additional side effects found with the individual drugs in this treatment. It does not list all possible side effects. For more information, talk with your healthcare provider.

Common Side Effects	Less Common Side Effects	Rare Side Effects
<ul style="list-style-type: none"> Fatigue or weakness 	<ul style="list-style-type: none"> Headache Increased tearing and eye irritation Muscle or joint pain Dizziness Difficulty sleeping Changes in taste 	<ul style="list-style-type: none"> Blood clots in the lungs or legs and related conditions, such as stroke Chest pain or heart failure Dehydration

Notes

Disclaimer. The drug information presented is time-sensitive and current as of the date stated below. It is not a substitute for the expertise, skill, knowledge and judgment of healthcare practitioners. Additional, important information about this product, including “black box” warnings and precautions, is available from the manufacturer and other sources. ONMARK PROVIDES THIS DRUG INFORMATION ON AN “AS IS” BASIS. ALL WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE ARE DISCLAIMED.

Limitation of Liability. Onmark shall not be liable for any improper or incorrect use of the information described and contained herein and assumes no responsibility for anyone’s use of the information. IN NO EVENT SHALL ONMARK BE LIABLE FOR ANY DIRECT, INDIRECT, INCIDENTAL, SPECIAL, EXEMPLARY, OR CONSEQUENTIAL DAMAGES (INCLUDING, BUT NOT LIMITED TO, PROCUREMENT OR SUBSTITUTE GOODS OR SERVICES; LOSS OF USE, DATA OR PROFITS; OR BUSINESS INTERRUPTION) HOWEVER CAUSED AND ON ANY THEORY OF LIABILITY, WHETHER IN CONTRACT, STRICT LIABILITY, OR TORT (INCLUDING NEGLIGENCE OR OTHERWISE) ARISING IN ANY WAY OUT OF THE USE OF THIS INFORMATION, EVEN IF ADVISED OF THE POSSIBILITY OF SUCH DAMAGE.

This document is a complimentary service provided by Onmark for its members. www.onmarkservices.com
 © 2008 Onmark, Inc. Onmark is a trademark of Oncology Therapeutics Network

January 2008

References:

- Fernando et al. A phase II study of oxaliplatin, capecitabine and bevacizumab in the treatment of metastatic colorectal cancer. Proc Am Soc Clin Oncol. 2005; Abstract 3556.
- Schmiegel et al. Comparable safety and response rate with bevacizumab in combination with capecitabine/oxaliplatin (CapOx/Bev) versus capecitabine/irinotecan (CapIri/Bev) in advanced CRC (mCRC): A randomized phase II study of the AIO GI tumor study group. Proc Am Soc Clin Oncol. 2007; Abstract 4034.

