

Patient Treatment Information

CISPLATIN IP + CYCLOPHOSPHAMIDE

Your chemotherapy treatment is called cisplatin IP + cyclophosphamide. It is commonly used to treat ovarian cancer and has also been used to treat other diseases. Cisplatin IP + cyclophosphamide is made up of two chemotherapy drugs:

- cisplatin (sis-PLA-tin) or Platinol (PLA-tin-all)
- cyclophosphamide (sy-e-kloe-FOSS-fa-mide)

These drugs prevent cancer cells from dividing and growing, and can eventually cause the cancer cells to shrink and die.

What Do I Need to Know Before Starting Chemotherapy?

Be sure to tell your healthcare provider about any prescription or over-the-counter products you are taking, including dietary supplements, vitamins, herbal medicines and homeopathic remedies.

Use an effective birth control method while you are taking these drugs. Chemotherapy drugs can cause harm to a fetus, so be sure to tell your healthcare provider right away if you or your partner become pregnant.

Avoid breastfeeding during treatment. It is not known if these drugs pass into breast milk.

Some chemotherapy drugs can cause sterility. Talk with your healthcare provider about your options if you want to have children in the future.

Do not get any immunizations or vaccinations while taking chemotherapy drugs without the approval of your healthcare provider.

What Do I Need to Know Before Starting This Treatment?

Cisplatin can cause an allergic reaction. In rare cases, the reaction can be severe and life-threatening. Tell your healthcare provider right away if you have trouble breathing, sudden swelling, hives or rash during your treatment.

Kidney damage is a rare side effect of cisplatin. The risk of this side effect increases with higher doses and the longer you are treated with the drug. Your healthcare provider will give you fluids before your treatment and may ask you to drink extra fluids after your treatment to protect your kidneys.

Damage to the ear is a rare side effect of cisplatin. The risk of this side effect increases with higher doses and the longer you are treated with the drug. Symptoms include ringing in the ears, hearing loss, dizziness and vertigo. The hearing loss first occurs in the high frequency range. Decreased ability to hear normal conversation can also occur. Tell your healthcare provider if you develop ringing in the ears, hearing loss in one or both ears, dizziness or vertigo.

Intraperitoneal instillation of cisplatin commonly causes abdominal pain. The pain responds to pain medicines and usually resolves within 24 hours.

Cyclophosphamide can cause delayed wound healing.

Cyclophosphamide can cause bleeding in the bladder. Talk with your healthcare provider about drinking extra fluids and emptying your bladder frequently to reduce the risk of this side effect. Call your healthcare provider right away if you have painful, frequent, or difficult urination, or blood in the urine.

Your treatment can interact with other medicines, including:

- Digoxin (Lanoxin)
- Cimetidine (Tagamet)
- Drugs used for seizures, such as phenytoin (Dilantin), valproic acid and carbamazepine (Tegretol)
- Furosemide (Lasix)

You should not take this treatment if you are allergic to cisplatin, cyclophosphamide or any components of these drugs

How Is the Treatment Given?

Your healthcare provider will give you cyclophosphamide by injection into a vein. Cisplatin will be instilled into the peritoneal cavity, the area that contains the abdominal organs. The dose you receive will be based on your weight and height. Your healthcare provider will determine the number of treatments you receive.

You may be given medicines to help prevent and control nausea and vomiting before you receive your treatment. These medicines may be given either by mouth or by injection into a vein.

What Are the Important Side Effects?

All drugs can cause side effects, but every person reacts differently to each drug. The following chart lists important side effects that can occur with your treatment, how often the side effect occurs (common, less common or rare), how to recognize and minimize side effects and possible treatments. Call your healthcare provider if you have any questions or concerns about side effects.

Potential Side Effects	How to Minimize Side Effects	Possible Treatments
Infection (Common) <ul style="list-style-type: none"> • Fever and chills • Painful urination • Sore throat and cough • Nasal congestion • Swelling or redness of the skin at the site of a wound 	<ul style="list-style-type: none"> • Wash your hands often. • Brush and floss your teeth daily. • Clean cuts right away with warm water, soap and antiseptic. • When your white blood cell count is low, stay away from crowds and people with colds or other illnesses. 	<ul style="list-style-type: none"> • You may be given medicine to increase your white blood cell count. • You may be given an antibiotic to treat or prevent infection. • Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.
Anemia (Common) <ul style="list-style-type: none"> • Fatigue or weakness • Dizziness • Pale skin • Feeling out of breath • Feeling cold 	<ul style="list-style-type: none"> • Plan rest periods throughout the day. • Organize daily activities so that you conserve your energy. • Try to eat a well balanced diet and drink plenty of fluids. • Stand up slowly to avoid getting dizzy. 	<ul style="list-style-type: none"> • You may be given medicine to increase your red blood cell count. • Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.
Nausea/Vomiting (Common) <ul style="list-style-type: none"> • Feeling queasy or sick to your stomach 	<ul style="list-style-type: none"> • Eat small, frequent meals and bland foods—such as bananas, rice, applesauce and toast. • Eat food cold or at room temperature so the smell of food will not bother you. • Avoid fried, spicy or fatty foods. • Eat and drink slowly. • Drink plenty of liquids during the day, but to avoid bloating, do not drink during meals. 	<ul style="list-style-type: none"> • You will be given medicine to help reduce nausea and vomiting.

Potential Side Effects	How to Minimize Side Effects	Possible Treatments
<p>Anorexia or Appetite Loss (Common)</p> <ul style="list-style-type: none"> • Not having an appetite • Feeling too nauseous to eat • Metallic or medicinal taste • Change in taste causing dislike for certain foods 	<ul style="list-style-type: none"> • Try eating six to eight small meals or snacks each day instead of three larger meals. • Vary your diet and try new foods and recipes. • Take a walk before meals, when possible. This may make you feel hungrier. • Eat with friends or family. When eating alone, listen to the radio or watch TV. 	<ul style="list-style-type: none"> • Cook dinners ahead of time and freeze them in small portions so that cooking smells are minimized. • Let others help with food, but ask that foods be prepared in small portions that can be frozen. And don't hesitate to let them know which foods to avoid. • Add mild spices to change flavor. • It might be helpful to have a program, such as Meals on Wheels, deliver food to you.
<p>Alopecia or Hair Loss (Common)</p> <ul style="list-style-type: none"> • More than normal amount of hair loss in your brush, in the shower or on your pillow after sleeping • Loss of body hair 	<p>Alopecia cannot be prevented but here are tips to help with hair loss:</p> <ul style="list-style-type: none"> • Use a soft hairbrush. Do not use brush rollers, color treat your hair or get a permanent. • Avoid daily hair washing, use a mild shampoo and avoid using a hairdryer, or use a low setting if you must use one. • Have your hair cut short; this will make it look fuller. • Your insurance might cover a wig. If you would like a wig, ask your healthcare provider for a prescription for a "hair prosthesis". Your hair color and style can be better matched if you shop for a wig before losing a lot of hair. • Use sunscreen or wear a hat or scarf to protect your scalp from the sun. 	
<p>Diarrhea (Less Common)</p> <ul style="list-style-type: none"> • Loose or watery stools several times a day • Abdominal cramping, gas and bloating 	<ul style="list-style-type: none"> • Eat small, frequent meals and bland foods—such as bananas, rice, applesauce and toast. • Avoid caffeine; alcohol; raw fruits and vegetables; raw eggs; undercooked meats; spicy, fatty and greasy foods; milk and dairy products; foods that cause gas, such as beans and other legumes; high fiber and high-fat foods; foods left un-refrigerated for more than two hours (one hour for egg dishes and cream or mayonnaise-based foods); bulk laxatives; and stool softeners. 	<ul style="list-style-type: none"> • Drink eight to ten glasses of clear liquids every day. • Your healthcare provider may prescribe medicine to help treat diarrhea.
<p>Mouth Sores and Pain (Less Common)</p> <ul style="list-style-type: none"> • Pain, swelling and redness of the mouth, tongue and throat • "Coated tongue" • Difficulty talking, swallowing or eating • Bleeding ulcers and infection 	<ul style="list-style-type: none"> • Brush teeth two to four times a day using a soft bristle brush and fluoride toothpaste. • Use non-waxed dental floss daily. • Ask your healthcare provider to recommend a mouthwash that does not contain alcohol. • Sip water during the day and use sugar-free candy or gum to keep your mouth wet. • Eat food cold or at room temperature. • Eat soft or pureed food. • Avoid food that is acidic, spicy, salty, dry or rough, such as toast. 	<ul style="list-style-type: none"> • You may be given medicine to help treat pain. • You may be given medicine to treat fungal or viral infections.
<p>Bleeding (Rare)</p> <ul style="list-style-type: none"> • Unusual bleeding, easy bruising • Black or tar-like stools • Blood in your urine • Pinpoint red spots on your skin • Bleeding gums or nosebleeds 	<ul style="list-style-type: none"> • Avoid aspirin and aspirin-like drugs, such as ibuprofen. • Use caution with sharp objects, such as razors and nail cutters. • Avoid activities that can cause cuts, bumps and bruises. 	<ul style="list-style-type: none"> • You may be given medicine to increase your platelet count. • Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.

Are There Other Possible Side Effects?

The following chart lists additional side effects found with this treatment. It does not list all possible side effects. For more information, talk with your healthcare provider.

Common Side Effects	Less Common Side Effects	Rare Side Effects
<ul style="list-style-type: none"> • Loss of balance • Skin discoloration • Darkening of nail beds • Stopping of menstrual periods in women or premature menopause 	<ul style="list-style-type: none"> • Metallic taste or loss of taste • Trouble walking • Congestion or sneezing • Fatigue • Eye irritation 	<ul style="list-style-type: none"> • Rash • Chest pain, heart attack, stroke • Shortness of breath • Lung problems, scarring of lung tissue • Secondary cancer or leukemia

When Should I Contact My Healthcare Provider?

Contact your healthcare provider if you have any of the following symptoms:

- Vomiting that is severe or lasts several hours
- Diarrhea that causes an additional four bowel movements a day, diarrhea at night or diarrhea with fever, cramps or bloody stools
- Shaking chills or fever higher than 100.5 degrees F
- Painful or frequent urination or blood in your urine
- Unusual bleeding, easy bruising or pinpoint red spots on your skin
- Inability to eat or weight loss
- Irregular or rapid heart beat
- Chest pain, chest tightness or shortness of breath
- Dizziness or feeling light headed
- Pain, redness, swelling or sores in your mouth or throat that interfere with eating
- Ringing in the ears or hearing loss in one or both ears

Notes

Disclaimer. The drug information presented is time-sensitive and current as of the date stated below. It is not a substitute for the expertise, skill, knowledge and judgment of healthcare practitioners. Additional, important information about this product, including “black box” warnings and precautions, is available from the manufacturer and other sources. ONMARK PROVIDES THIS DRUG INFORMATION ON AN “AS IS” BASIS. ALL WARRANTIES OF ANY KIND, EXPRESS OR IMPLIED, INCLUDING BUT NOT LIMITED TO THE IMPLIED WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR PURPOSE ARE DISCLAIMED.

Limitation of Liability. Onmark shall not be liable for any improper or incorrect use of the information described and contained herein and assumes no responsibility for anyone’s use of the information. IN NO EVENT SHALL ONMARK BE LIABLE FOR ANY DIRECT, INDIRECT, INCIDENTAL, SPECIAL, EXEMPLARY, OR CONSEQUENTIAL DAMAGES (INCLUDING, BUT NOT LIMITED TO, PROCUREMENT OR SUBSTITUTE GOODS OR SERVICES; LOSS OF USE, DATA OR PROFITS; OR BUSINESS INTERRUPTION) HOWEVER CAUSED AND ON ANY THEORY OF LIABILITY, WHETHER IN CONTRACT, STRICT LIABILITY, OR TORT (INCLUDING NEGLIGENCE OR OTHERWISE) ARISING IN ANY WAY OUT OF THE USE OF THIS INFORMATION, EVEN IF ADVISED OF THE POSSIBILITY OF SUCH DAMAGE.

This document is a complimentary service provided by Onmark for its members. www.onmarkservices.com
 © 2006 Onmark, Inc. Onmark is a trademark of Oncology Therapeutics Network

November 2006

