

Your chemotherapy treatment is called FAC. It is commonly used to treat breast cancer and has also been used to treat other diseases. FAC is made up of three chemotherapy drugs:

- 5-fluorouracil (5-FU), (five-flure-oh-YOOR-a-sill)
- doxorubicin (dox-oh-ROO-bi-sin) or Adriamycin® (a-dree-a-MY-sin)
- cyclophosphamide (sy-e-kloe-FOSS-fa-mide) or Cytosan® (sy-e-TOX-an)

These drugs prevent cancer cells from dividing and growing, and can eventually cause the cancer cells to shrink and die.

## **What Do I Need to Know Before Starting Treatment?**

Be sure to tell your healthcare provider about any prescription or over-the-counter products you are taking, including dietary supplements, vitamins, herbal medicines and homeopathic remedies.

Use an effective birth control method while you are taking these drugs. Chemotherapy drugs can cause harm to a fetus, so be sure to tell your healthcare provider right away if you or your partner become pregnant.

Avoid breastfeeding during treatment. It is not known if these drugs pass into breast milk.

Some chemotherapy drugs can cause sterility. Talk with your healthcare provider about your options if you want to have children in the future.

Do not get any immunizations or vaccinations while taking chemotherapy drugs without the approval of your healthcare provider.

## **What Do I Need to Know Before Starting FAC?**

Doxorubicin can cause heart problems, such as congestive heart failure, during treatment or months or years after your treatment ends. Your risk of heart damage from doxorubicin increases as the total dose of doxorubicin or similar chemotherapy medicines you take increases. 5-FU can cause heart problems, including an irregular heart beat, angina, a heart attack or heart failure. These side effects may be more common in patients with a history of heart disease. Tell your healthcare provider if you have been treated with chemotherapy, radiation therapy to the chest or if you have heart disease. Call your healthcare provider immediately if you have back pain, flushing or chest tightness.

Doxorubicin can cause irritation if it accidentally leaks out of the vein. Tell your healthcare provider right away if you feel pain, burning or swelling around the injection site during your treatment.

Doxorubicin can cause urine, tears or sweat to turn red for one to two days after treatment. The red color is caused by the medicine and is not harmful.

In rare cases, doxorubicin can cause secondary cancers or leukemia years after treatment.

Cyclophosphamide can cause delayed wound healing.

Cyclophosphamide can cause bleeding in the bladder. Talk with your healthcare provider about drinking extra fluids and emptying your bladder frequently to reduce the risk of this side effect. Call your healthcare provider right away if you have blood in the urine or painful, frequent or difficult urination.

Your treatment can interfere with other medicines, including:

- Cimetidine (Tagamet®)
- Digoxin (Lanoxin®) or verapamil
- Phenytoin (Dilantin®) or phenobarbital
- Cyclosporine (Sandimmune®)

Please note this list is a summary and does not contain all possible drug interactions. Contact your healthcare provider if you are taking any medications that can interact with your treatment.

**You should not take this treatment if you:**

- Are allergic to 5FU, doxorubicin, cyclophosphamide or any components of these drugs.
- Have a condition known as dihydropyrimidine dehydrogenase (DPD) deficiency.

### How Is the Treatment Given?

Your healthcare provider will give you your treatment by injection into a vein. The dose you receive will be based on your weight and height. Your healthcare provider will determine the number of treatments you receive.

You may be given medicines to help prevent and control nausea and vomiting before you receive your treatment. These medicines may be given either by mouth or by injection into a vein.

If you are given any medicine to take at home, do not share it with others. Sharing this medication with anyone else could be harmful.

### When Should I Call My Healthcare Provider?

Call your healthcare provider right away if you have any of the following symptoms:

- Shaking chills or fever of 100.5 degrees F or higher
- Unusual bleeding, easy bruising or pinpoint red spots on your skin
- Vomiting that is severe or lasts several hours
- Painful or frequent urination or blood in your urine
- Diarrhea that causes an additional four bowel movements a day, diarrhea that lasts more than one day, diarrhea at night or diarrhea with fever, cramps or bloody stools
- Irregular or rapid heart beat, chest pain, chest tightness or shortness of breath
- Dizziness or feeling lightheaded
- Inability to eat or weight loss

### What Are the Possible Side Effects?

All drugs can cause side effects, but every person reacts differently to each drug. The following chart lists the possible side effects that can occur with your treatment, how to recognize and minimize symptoms and possible treatments. The side effects are grouped by how often the side effect occurs: Common (occurs in more than 25 percent of patients), Less Common (occurs in 5 to 25 percent of patients) or Rare (occurs in less than 5 percent of patients).

Side Effect	How to Minimize Side Effect	Possible Treatments
<b>Risk of Infection (Common)</b> <ul style="list-style-type: none"><li>• Fever and chills</li><li>• Painful urination</li><li>• Sore throat and cough</li><li>• Nasal congestion</li><li>• Swelling or redness of the skin at the site of a wound</li></ul>	<ul style="list-style-type: none"><li>• Wash your hands often.</li><li>• Brush and floss your teeth daily.</li><li>• Clean cuts right away with warm water, soap and antiseptic.</li><li>• When your white blood cell count is low, stay away from crowds and people with colds or other illnesses.</li></ul>	<ul style="list-style-type: none"><li>• You may be given medicine to increase your white blood cell count.</li><li>• You may be given an antibiotic to treat or prevent infection.</li><li>• Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.</li></ul>
<b>Nausea/Vomiting (Common. Symptoms are generally mild.)</b> <ul style="list-style-type: none"><li>• Feeling queasy or sick to your stomach</li></ul>	<ul style="list-style-type: none"><li>• Eat small, frequent meals and bland foods—such as bananas, rice, applesauce and toast.</li><li>• Eat food cold or at room temperature so the smell of food will not bother you.</li><li>• Avoid fried, spicy or fatty foods.</li><li>• Eat and drink slowly.</li><li>• Drink plenty of liquids during the day, but to avoid bloating, drink small amounts of liquid during meals.</li></ul>	<ul style="list-style-type: none"><li>• You will be given medicine to help reduce nausea and vomiting.</li></ul>
<b>Anemia (Less Common)</b> <ul style="list-style-type: none"><li>• Fatigue or weakness</li><li>• Dizziness</li></ul>	<ul style="list-style-type: none"><li>• Plan rest periods throughout the day.</li><li>• Organize daily activities so that you conserve your energy.</li></ul>	<ul style="list-style-type: none"><li>• You may be given medicine to increase your red blood cell count.</li><li>• Your healthcare provider may decrease</li></ul>

Side Effect	How to Minimize Side Effect	Possible Treatments
<ul style="list-style-type: none"> <li>• Pale skin</li> <li>• Feeling out of breath</li> <li>• Feeling cold</li> </ul>	<ul style="list-style-type: none"> <li>• Try to eat a well balanced diet and drink plenty of fluids.</li> <li>• Stand up slowly to avoid getting dizzy.</li> </ul>	<ul style="list-style-type: none"> <li>• your chemotherapy dose or delay further chemotherapy.</li> </ul>
<p><b>Mouth Sores and Pain (Less Common. Symptoms are generally mild.)</b></p> <ul style="list-style-type: none"> <li>• Pain, swelling or redness of the mouth, tongue and throat</li> <li>• “Coated tongue”</li> <li>• Difficulty talking, swallowing or eating</li> <li>• Bleeding ulcers and infection</li> </ul>	<ul style="list-style-type: none"> <li>• Brush teeth two to four times a day using a soft bristle brush and fluoride toothpaste.</li> <li>• Use non-waxed dental floss daily.</li> <li>• Ask your healthcare provider to recommend a mouthwash that does not contain alcohol.</li> <li>• Sip water during the day and use sugar-free candy or gum to keep your mouth wet.</li> <li>• Eat food cold or at room temperature.</li> <li>• Eat soft or pureed food.</li> <li>• Avoid food that is acidic, spicy, salty, dry or rough, such as toast.</li> <li>• Sucking on ice chips for 10 to 15 minutes before and after fluorouracil injections may help relieve the pain of mouth sores.</li> </ul>	<ul style="list-style-type: none"> <li>• You may be given medicine to help treat pain.</li> <li>• You may be given medicine to treat fungal or viral infections.</li> </ul>
<p><b>Diarrhea (Less Common. Symptoms are generally mild.)</b></p> <ul style="list-style-type: none"> <li>• Loose or watery stools several times a day</li> <li>• Abdominal cramping, gas and bloating</li> </ul>	<ul style="list-style-type: none"> <li>• Eat small, frequent meals and bland foods—such as bananas, rice, applesauce and toast.</li> <li>• Avoid caffeine; alcohol; raw fruits and vegetables; raw eggs; undercooked meats; spicy, fatty and greasy foods; milk and dairy products; foods that cause gas, such as beans and other legumes; high fiber and high-fat foods; foods left un-refrigerated for more than two hours (one hour for egg dishes and cream or mayonnaise-based foods); bulk laxatives; and stool softeners.</li> </ul>	<ul style="list-style-type: none"> <li>• Drink eight to ten glasses of clear liquids every day.</li> <li>• Your healthcare provider may prescribe medicine to help treat diarrhea.</li> </ul>
<p><b>Bleeding (Rare)</b></p> <ul style="list-style-type: none"> <li>• Unusual bleeding or easy bruising</li> <li>• Black or tar-like stools</li> <li>• Blood in your urine</li> <li>• Pinpoint red spots on your skin</li> <li>• Bleeding gums or nosebleeds</li> </ul>	<ul style="list-style-type: none"> <li>• Avoid aspirin and aspirin-like drugs, such as ibuprofen.</li> <li>• Use caution with sharp objects, such as razors and nail cutters.</li> <li>• Avoid activities that can cause cuts, bumps and bruises.</li> </ul>	<ul style="list-style-type: none"> <li>• You may be given medicine to increase your platelet count.</li> <li>• Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.</li> </ul>
<p><b>Neuropathy (Rare. Symptoms are generally mild.)</b></p> <ul style="list-style-type: none"> <li>• Numbness or tingling feeling in the hands or feet</li> <li>• Muscle cramps</li> <li>• Loss of balance</li> <li>• Difficulty buttoning buttons or picking up objects</li> <li>• Decreased awareness of heat or cold in fingertips and toes</li> <li>• Difficulty hearing</li> </ul>	<ul style="list-style-type: none"> <li>• Try to avoid the cold or extreme heat.</li> <li>• Wear mittens or gloves, socks and scarves.</li> <li>• If your fingers are numb, be careful with sharp objects.</li> <li>• Beware of hot coffee mugs, pots and pans and dishwasher—you may not feel the heat until you are burned.</li> <li>• If you feel unsteady, be careful on stairs and in the shower.</li> </ul>	<ul style="list-style-type: none"> <li>• Your healthcare provider may decrease your chemotherapy dose or delay further chemotherapy.</li> </ul>
<p><b>Radiation Recall (Rare)</b></p> <ul style="list-style-type: none"> <li>• Redness, tenderness or swelling on areas of the skin that have previously been treated with radiation.</li> <li>• May include wet sores, peeling skin or discoloration after the skin has healed.</li> </ul>	<ul style="list-style-type: none"> <li>• Stay out of the sun and avoid tanning beds.</li> <li>• If you are in the sun, wear protective clothing and use sunscreen with an SPF of 30 or higher.</li> </ul>	<ul style="list-style-type: none"> <li>• You may be given a corticosteroid to reduce swelling.</li> </ul>

